

Efficient frequency calculations in QM/MM using the Mobile Block Hessian method

An Ghysels¹, H. Lee Woodcock III², Yihan Shao³, Jing Kong³, Bernard R. Brooks⁴,
Dimitri Van Neck¹, Veronique Van Speybroeck¹, Michel Waroquier¹

¹ Center for Molecular Modeling, Ghent University, Ghent, Belgium

² Department of Chemistry, University of South Florida, Tampa, FL

³ Q-Chem Inc., Pittsburgh, PA

⁴ Laboratory of Computational Biology, NHLBI, NIH, Bethesda, MD

Normal mode analysis (NMA) is a well-known technique which estimates the intrinsic frequencies of chemical systems by assuming a harmonic shape for the potential energy surface. Despite its simplicity, it is still a popular approach to predict vibrational IR and Raman spectra, to identify chemical groups [1], or to study the large motions involved in conformational changes of biomolecules [2]. The method is based on the diagonalization of the Hessian matrix, which contains the second derivatives of the potential energy with respect to the $3N$ nuclear coordinates. In extended QM/MM systems, however, its calculation, storage and diagonalization is an expensive computational task. Even in the case of a small QM region, the numerous derivatives of the QM/MM interaction terms still form a bottleneck in the frequency calculation: each MM atom displacement affects the charge distribution in the QM region, resulting in a large number of coupled perturbation self-consistent field (CPSCF) equations [3].

Recently, the Mobile Block Hessian (MBH) method was developed in order to reduce the dimensionality of the Hessian [4,5]. The main concept is the introduction of blocks that move as rigid bodies during the vibrational analysis. This block concept is now combined with the QM/MM scheme. Block motions replace the individual atom motions in the CPSCF equations, thus reducing the number of equations drastically.

We have derived and implemented a parallel version of both the full QM/MM Hessian calculation and the QM/MM Mobile Block Hessian calculation in the Q-Chem/CHARMM [6,7] interface. The reduced computational cost opens the path to a broad range of applications of normal mode analysis. As an example, the oxidative deboronation of the drug bortezomib is presented, which is suspected to induce cell death of cancer cells [8]. Application of MBH in the QM/MM scheme realizes a speed up of a factor 200 compared to the full QM calculation, while the estimated free energy difference are within less than 1 kJ/mol accuracy.

[1] R. J. Fessenden and J. S. Fessenden. Organic chemistry, 4th ed., 1990.

[2] Q. Cui and I. Bahar. Normal Mode Analysis: Theory and applications to biological and chemical systems, 2006.

[3] Q. Cui and M. Karplus. J. Chem. Phys. 112, 1133-1149, 2000.

[4] A. Ghysels, D. Van Neck, V. Van Speybroeck, T. Verstraelen, and M. Waroquier. J. Chem. Phys. 126, 224102, 2007

[5] A. Ghysels, D. Van Neck, V. Van Speybroeck, B. R. Brooks, and M. Waroquier. J. Chem. Phys. 130, 084107, 2009

[6] Y. Shao et al. Phys. Chem. Chem. Phys. 8(27), 3172-3191, 2006.

[7] B. R. Brooks et al. J. Comput. Chem. 30(10), 1545-1614, 2009.

[8] J. D. Larkin, K. L. Bhat, G. D. Markham, T. D. James, B. R. Brooks, C. W. Bock, J. Phys. Chem. A 112, 8446-8454, 2008.