

---

## Modeling aldol condensations in UiO-66 type materials

J. Hajek<sup>1</sup>, M. Vandichel<sup>1</sup>, F. Vermoortele<sup>2</sup>, M. Waroquier<sup>1</sup>, D. De Vos<sup>2</sup>, V. Van Speybroeck<sup>1</sup>

1. Center for Molecular Modeling, Ghent University, Technologiepark 903, B-9052 Zwijnaarde, Belgium
2. Centre for Surface Chemistry and Catalysis, University of Leuven, Kasteelpark Arenberg 23, B-3001 Leuven, Belgium

Hybrid framework compounds including both inorganic and organic moieties have recently developed into an important new class of crystalline porous materials, also known as metal organic frameworks (MOFs) [1]. MOFs contain many metal sites, which might be catalytically active. This implies that MOFs have the potential to replace homogenous catalysts in important industrial applications [2]. Notwithstanding the catalytic activity of some of them- it is not always clear what the active sites are and how the reactions are catalyzed. In this contribution, we will focus on certain variations of the UiO-66. Recently, the UiO-66-NH<sub>2</sub> has successfully been applied experimentally for the synthesis of Jasminaldehyde from benzaldehyde and heptanal [3]. To fully understand the mechanism of Jasminaldehyde condensation on such open active Zr-sites, we applied molecular modeling techniques on extended cluster as well as on periodic models of the UiO-66-X materials. On an extended cluster, the most plausible condensation reaction mechanism was found, showing that benzaldehyde and heptanal are adsorbed and activated on adjacent Zr sites before the coupling reaction. On the UiO-66-NH<sub>2</sub> model, there should be also an alternative reaction route, where one of the two aldehydes is activated on the amino group, explaining the higher selectivity of the UiO-66-NH<sub>2</sub> compared to the UiO-66 [3]. Deeper insight into these complex mechanisms allows us to explain the selectivity and activity on the UiO-66 type materials.

[1] Férey, G., Chemical Society Reviews 37 (2008) 191.

[2] Czaja, A. U., Trukhan, N. and Muller, U., Chem. Soc. Rev. 38 (2009) 1284.

[3] Vermoortele, F., Ameloot, R., Vimont, A., Serre, C., De Vos, D., Chemical Communications 47 (2011) 1521.