

Selectivity prediction: making the comparison between extended cluster and periodic calculations

Matthias Vandichel,¹ Frederik Vermoortele,² Stijn Cottenie,¹ Dirk De Vos,²

Michel Waroquier¹ and Veronique Van Speybroeck¹

¹Center for Molecular Modeling, Technologiepark 903, 9052 Zwijnaarde, Belgium

²Centre for Surface Chemistry and Catalysis, Kasteelpark Arenberg 23, 3001 Leuven, Belgium

Abstract

In this work, the catalytic properties of the metal organic framework (MOF) Cu₃BTC₂ have been studied as catalyst. More in particular, the selectivity for the isopulegol isomers as products of the citronellal cyclization has been theoretically investigated (Fig. 1). Recently, this reaction was also studied to investigate the electronic effects of linker substitution on the lewis acid MOF UiO-66 [1].




Figure 1. Four possible isomers are formed from the citronellal cyclization

To find a plausible model for the citronellal cyclization (Fig. 1), we performed extended cluster and periodic calculations (Fig. 2a-b) within DFT. All relevant reaction routes leading to the various isopulegol isomers are determined theoretically for each catalyst model. We used Curtin Hammett's principle, stating that only Gibbs free energy differences between the transition states are necessary to predict initial selectivities. Those selectivities could then be compared with experimental values obtained on highly pure Cu₃BTC₂ crystals [2]. In this work, we focus on the selectivity towards isopulegol. For the extended cluster calculations, the transition states were optimized using an ONIOM-approach, with energy refinements at B3LYP/6-311+g(d,p)-D3, while the periodic calculations were performed at the PBE-D2 level, applying very robust optimization parameters.

The experimental selectivity towards isopulegol drops when temperature raises. This general trend is reproduced by both theoretical approaches (Fig. 3). Some experimental data points at

low citronellal conversion are also shown in Fig. 3. The selectivity in the periodic approach agrees well with the experimental behavior.




Figure 2. Cyclization of citronellal towards isopulegol: (a) extended cluster, versus (b) periodic calculations




Figure 3. Isopulegol selectivity in function of temperature

Summarizing, relevant transition states are determined to have an accurate description of the citronellal cyclization on Cu_3BTC_2 . The periodic calculations comprise a complete description of the catalyst environment by construction, which can not be fully covered by the extended cluster model. This might explain why only periodic PBE-D2 calculations are able to accurately predict the selectivity at various reaction temperatures.

Keywords: periodic DFT-D, selectivity prediction, citronellal cyclization

- [1] Vermoortele, F., Vandichel, M., Van de Voorde, B., Ameloot, R., Waroquier, M., Van Speybroeck, V., De Vos, D. E., *Angewandte Chemie International Edition*, in press
[2] Vandichel, M., Vermoortele, F., Cottenie, S., Devos, D.E., Waroquier, M., Van Speybroeck, V., submitted